


Mapping America – Number 129

“Has Ever Had an STD” by Marital Status and Religious Attendance

Percentage Who Have Ever Had an STD
by Current Religious Attendance (adults aged 18-59)


Source: National Health and Social Life Survey, 1992



Althea Nagai, Ph.D.

Percentage Who Have Ever Had an STD
by Marital Status (adults aged 18-59)


Source: National Health and Social Life Survey, 1992



Althea Nagai, Ph.D.

Percentage Who Have Ever Had an STD
by Marital Status and Religious Attendance (adults aged 18-59) Combined

Source: National Health and Social Life Survey, 1992



Althea Nagai, Ph.D.

This chart looks at the number of adults who have ever had an STD at the extremes of four demographic quadrants.

These four quadrants are derived from combining two sets of marital statuses (always-intact vs. non-intact* and singles) and two sets of religious attendance (high vs. low).

The individuals occupying the four corners (or four extremes) of these quadrants are:

The always-intact married individual that worships weekly;

The always-intact married individual that never worships;

The non-intact* or single individual that worships weekly;

The non-intact* or single individual that never worships.

(* The non-intact group consists of divorced or separated and divorced and remarried individuals.)

Has Ever Had an STD

Adults aged 18 to 59 in intact marriages and adults who worship weekly were least likely to have ever had a sexually transmitted disease, according to the National Health and Social Life Survey (1992), the most detailed analysis of sexual behavior in America.

Religious attendance: Those who worshiped weekly were least likely to have ever had an STD (1.3 percent). The likelihood of having an STD increased as regularity of worship decreased: 2 percent of those who worshiped less than monthly have ever had an STD; 2.5 percent of those who worshiped less than weekly but at least monthly had ever had an STD and 2.8 percent of those who never worshiped had ever had an STD.

Marital status: Those in always-intact marriages were least likely to have ever had a sexually transmitted disease (1.3 percent). Sexually transmitted disease is more prevalent in non-intact family structures and among singles: 1.8 percent of those who were always single have had a sexually transmitted disease; 3 percent of those who were divorced or separated have had a sexually transmitted disease and 3.1 percent of those who were divorced and remarried had ever had a sexually transmitted disease.

Religious attendance and marital status combined: Those in non-intact family structures or who were single who worshiped weekly were less likely to have ever had an STD (1.1 percent), followed by those in intact marriages who worshiped weekly (1.6 percent). Those in intact marriages who never worshiped (2.7 percent) and those in non-intact family structures or who were single who never worshiped (3.1 percent) were more likely to have ever had an STD. (The healthiest score reported here is unusual.)

Related Insight from Other Studies

Analysis of Wave One of the National Longitudinal Study of Adolescent Health (Add Health) found that STD reports were generally higher among adolescents who were not living in two-parent homes.¹

Data from the 1992 Youth Risk Behavior Survey/Supplement to the National Health Interview Survey found that living in a two-parent family was protective against adolescent sexual intercourse as well as STD transmission.²

Another study shows that a history of STDs tends to coincide with a history of drinking problems.³

Patrick F. Fagan, Ph.D. and Althea Nagai, Ph.D.

Dr. Fagan is senior fellow and director of the Marriage and Religion Research Institute (MARRI) at Family Research Council.

Althea Nagai is a visiting fellow at Family Research Council.

-
- 1 E.C. Newbern, W.C. Miller, V.J. Schoenback, and J.S. Kaufman, "Family Socioeconomic Status and Self-Reported Sexually Transmitted Diseases among Black and White American Adolescents," *Journal of Sexually Transmitted Diseases* 31, no. 9 (September 2004): 533-541.
 - 2 J.S. Santelli, R. Lowry, N.D. Brener and L. Robin, "The Association of Sexual Behaviors with Socioeconomic Status, Family Structure, and Race/Ethnicity among US Adolescents," *Journal of Public Health* 90, no. 10 (October 2000): 1582-1588.
 - 3 Karen Paige Ericksen and Karen F. Trocki, "Sex, Alcohol and Sexually Transmitted Diseases: A National Survey," *Journal of Family Planning Perspectives* 26, no. 6 (November 1994): 257-263.

Marriage and Religion Research Institute / Editor: Patrick F. Fagan, Ph.D. / Managing Editors: Julia Kiewit and Anna Dorminey

Please feel free to make additional copies of this resource. We only ask that you comply with standard norms of attribution to the Family Research Council and to the Marriage and Religion Research Institute.

WWW.MAPPINGAMERICAPROJECT.ORG



MARRIAGE AND RELIGION
RESEARCH INSTITUTE
801 G STREET NW
WASHINGTON DC 20001
800-225-4008
WWW.MARRI.US

MA13E09